

Alberta Thy 20-99

October 1999

Holographic Counterterm Actions and Anomalies for Asymptotic AdS and Flat Spaces

Jeongwon Ho

Theoretical Physics Institute, Department of Physics,
University of Alberta, Edmonton, Canada T6G 2J1
E-mail: jwho@phys.ualberta.ca

Abstract

Counterterm actions are constructed along the ADM formalism. It is shown that the counterterm action can be intrinsically written in terms of intrinsic boundary geometry. Using the expression of counterterm action, we obtain a general form of the counterterm action available for any d -dimensional spherical boundary. In the description, we also derive *arbitrary* dimensional holographic conformal anomaly. It is also shown that counterterm actions for AF spaces can be obtained from the AdS description just as taking the limit of $\ell \rightarrow \infty$. An asymptotically flat spacetime with non-spherical boundary is speculated. In the example, additional counterterms to eliminate (leading) divergent terms due to deviation of boundary from round sphere are imagined by observing (4-dimensional) holographic anomaly proportional to $\square R$. Argument of the deceptive-like anomaly is given by comparing with the holographic description of 5-dimensional Kerr-AdS spacetime.

1 Introduction

There has been a typical problem to define a gravitational action suffering from divergence in a non-compact space. In spite that several prescriptions within the concept of reference space have been suggested so far [1][2][3], those are flawed by the fact although the divergences could be eliminated by choosing an appropriate reference space, it is impossible to embed a boundary with an arbitrary geometry. Another drawback of the reference space method is that different reference spaces are needed for different boundary geometries, so that one cannot define relative energies in a consistent manner.

Recently, a prominent prescription has been suggested [4] in the context of AdS/CFT correspondence [5][6][7], which could be understood as a realization of the holographic principle [8][9]. According to the correspondence, UV divergences of quantum field theory living on a boundary of AdS space are derived from IR divergences of the bulk theory (UV-IR connection [10]). So, the bulk action could be regularized by adding local counterterms [6][11]. On asymptotic AdS spaces, this approach gives an elegant expression of counterterm action in the form of the expansion for AdS radius ℓ [4][12][13][14]

$$\begin{aligned} \tilde{S} = & -\frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g_0} \left\{ \frac{d-1}{\ell} + \frac{\ell}{2(d-2)} R \right. \\ & \left. + \frac{\ell^3}{2(d-2)^2(d-4)} \left(R_{ab} R^{ab} - \frac{d}{4(d-1)} R^2 \right) + \cdots \right\}. \end{aligned} \quad (1)$$

In case of even dimensional boundary, however, one encounters a logarithmic divergent term in evaluating the bulk action functional. In order to obtain a finite action if we take the counterterm action involving this log term, it would cause problematic results in calculating a boundary stress energy tensor [13]. Even though the logarithmic divergence embarrasses to obtain a finite regularized action, it provides a remarkable consistency check of the AdS/CFT correspondence [6][11]. In other words, because a conformal anomaly for d -dimensional conformal field theory in coupling to background gravity comes from logarithmic UV divergences [15], evaluation of the conformal anomaly in this scheme becomes a nontrivial check of the AdS/CFT correspondence. (For holographic conformal anomaly for the dilaton coupled conformal field theory, see Ref.[16]).

The counterterm action of Eq.(1) has been also constructed from the Gauss-Codazzi equations through an iterative process [14]. In the Ref.[14], counterterm actions for asymptotically flat (AF) spaces have been also investigated. However, the procedure adapted for AdS spaces could not be simply generalized on AF descriptions because of mathematical difficulty due to non-linearity of the Gauss-Codazzi equations. Taking an alternative approach, they obtained a counterterm action for AF spaces with $S^{d-n} \times \mathbb{R}^n$ boundary geometries

$$\tilde{S} = -\frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g_0} \sqrt{\frac{R^3}{R^2 - R_{ab}R^{ab}}}. \quad (2)$$

Very recently, a different prescription to construct the counterterm action has been suggested in Ref.[17]. In the prescription, a length dimensional parameter analogue to the radius of AdS space was defined, so that the counterterm actions for asymptotically flat and AdS spaces are consistently constructed in the expansion for the new length parameter.

In this paper, we introduce another method to construct the counterterm actions in (1) and (2). In this construction, we take the ADM formalism and show that the counterterm action can be intrinsically written by the terms of intrinsic boundary geometry. Using our new expression for counterterm action, we obtain a general form of the counterterm action available for any d -dimensional spherical boundary. In the description, we also derive *arbitrary* dimensional holographic conformal anomaly. It is also shown that the counterterm action for AF spaces can be obtained from the AdS description just as taking the limit of $\ell \rightarrow \infty$.

On the other hand, a counterterm action for AF space with nontrivial boundary geometry is examined. Our counter example is the D -dimensional generalization of the Kerr metric [18] setting the mass parameter to zero. It is the metric of an AF space in spheroidal coordinates. This example has been considered in Ref.[14]. The authors have shown that for $d > 6$ the counterterm action in (2) based on round sphere boundary does not eliminate all divergent terms. In Ref.[17], it has been shown that leading order of divergent terms due to deviation from round sphere could be canceled by introducing an additional counterterm action whose a form is similar with the counterterms of squared boundary curvature in Eq.(1). Here, we shall show that the additional counterterms can be conjectured in somewhat interesting scheme

as well: Our derivation of the counterterm action does not use the full Einstein equations. Instead only normal-normal projection equation, which is obtained by projecting the Einstein equations on the boundary in normal directions, is used. In case of simple boundary geometry (round sphere), other equations, tangential-tangential and tangential-normal projections, are not crucial, because they become trivial or dummy on the procedure. However, in case of nontrivial boundary geometry, the full Einstein's equations must be used. Our observation is that taking only the normal-normal projection equation, we obtain additional divergent terms in boundary action value (BAV), which are logarithmic. Thus, the conformal invariance of regularized action (RA) would be broken by a conformal anomaly. However, we shall show that for an example of $d = 4$ the anomaly is proportional to $\square R$. That is, additional counterterms proportional to the squared boundary curvature can be added on the counterterm action in (2) and the conformal invariance would be recovered. We also briefly discuss about these aspects comparing with AdS descriptions.

Our paper is organized as follows; In Sect.2, counterterm action is constructed in the ADM formulation. Examples for asymptotic AdS spaces are considered in Sect.3; A counterterm action for asymptotic AdS space with S^d boundary is constructed. In this example, arbitrary dimensional conformal anomaly is obtained. In Sect.4, relationship between counterterm actions for asymptotic AdS and flat spaces is discussed. For an AF space in spheroidal coordinates, the divergent terms are evaluated. Additional counterterms to eliminate divergent terms due to deviation from round sphere are conjectured in observation of logarithmic divergence. We also give a brief discussion about similarities of these aspects and AdS descriptions in a holographic sense. Discussions and summary are contained in Sect.5.

2 Holographic Counterterm Actions

$(d + 1)$ -dimensional gravitational action with cosmological constant $\Lambda = -d(d - 1)/(2\ell^2)$ is given by

$$S = \frac{1}{16\pi G} \int_X d^{d+1}x \sqrt{-\hat{G}} \left(\hat{R} + \frac{d(d-1)}{\ell^2} \right) - \frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g} \Theta, \quad (3)$$

where g_{ab} is boundary metric and Θ is the trace of extrinsic curvature of d -dimensional timelike boundary ∂X defined by $\Theta_{ab} = -g_a^\mu \nabla_\mu n_b$. ∇ denotes the covariant derivative on $(d+1)$ -dimensional manifold X and n^μ is an outward unit normal to the boundary ∂X . The boundary term in Eq.(3), so called Gibbons-Hawking term, is required for well defined variational principle.

Our purpose is to add another proper surface integral to the action in (3), so that the action becomes finite in the limit that the boundary is taken to infinity. According to the counterterm subtraction approach, the additional surface integral must be written in terms of intrinsic boundary geometry. For the procedure, we take the ADM formulation as a guide line for construction of the counterterm action. As it will be seen in the following, the ADM formulation guarantees for the counterterm action to be written in terms of intrinsic boundary geometry.

To rewrite the action (3) in a canonical form, we first take a metric given by

$$\hat{G}_{\mu\nu} dx^\mu dx^\nu = N^2 d\rho^2 + g_{ab} dx^a dx^b, \quad (4)$$

where $N^2 = N^2(\rho)$ and $g = g(\rho, x^a)$. In this coordinate system, unit normal to the boundary is given by $n_\mu = N\delta_\mu^\rho$. Then, following the standard ADM procedure, canonical form of the action (3) becomes

$$S = \int_X d^{d+1}x (\pi^{ab} g'_{ab} - N\mathcal{H}_\rho) \equiv \int_X d^{d+1}x \mathcal{L}, \quad (5)$$

where $\pi^{ab} = \delta\mathcal{L}/\delta g'_{ab}$ is the momentum density conjugate to g_{ab} and $'$ denotes the derivative of ρ . ‘Hamiltonian’ density $\mathcal{H}_\rho$ is given by

$$\mathcal{H}_\rho = \frac{16\pi G}{\sqrt{-g}} \left(\frac{\pi^2}{d-1} - \pi_{ab}\pi^{ab} \right) - \frac{\sqrt{-g}}{16\pi G} \left(R + \frac{d(d-1)}{\ell^2} \right), \quad (6)$$

where R is d -dimensional scalar curvature of the boundary. The equation $\mathcal{H}_\rho = 0$ generates reparametrization of space coordinate ρ . In fact, this equation one of the Gauss-Codazzi equations that is defined by projecting the Einstein equations on the boundary in normal directions.

Using the constraint equation $\mathcal{H}_\rho = 0$, the BAV evaluated from the action (3) on the boundary $\rho = \rho_0$ is given by a simple form as

$$\begin{aligned} S_{cl} &= \int_{\partial X} d^d x \left\{ \frac{1}{8\pi G} \int^{\rho_0} d\rho N \sqrt{-g} \left(R + \frac{d(d-1)}{\ell^2} \right) \right\} \\ &\equiv \int_{\partial X} d^d x A(x^a; \rho_0). \end{aligned} \quad (7)$$

So, according to the counterterm subtraction approach, regularized action, S_{RA} , is defined by

$$S_{RA} \equiv S - \tilde{S}, \quad (8)$$

where the counterterm action $\tilde{S}$ is given by

$$\tilde{S} = - \int_{\partial X} d^d x \text{Div} (A(x^a; \rho_0)) \equiv - \int_{\partial X} d^d x \sqrt{-g_0} \bar{A}^{div}(x^a; \rho_0), \quad (9)$$

where Div means to pick divergent terms after ρ -integration and g_0 is the induced metric on the boundary.

What a counterterm action is a coordinate invariant functional of intrinsic boundary geometry is an important requirement in the counterterm subtraction method. This is because it is the only way to eliminate divergence of a gravitational action without disturbing the equations of motion or the symmetries [14]. In Eq.(9), the counterterm action (functional $A(x^a; \rho_0)$) is explicitly given in terms of intrinsic boundary geometry. (The ‘lapse’ function N can be absorbed in the space coordinate ρ by a coordinate redefinition.) In fact, the divergent terms in BAV is originated from the Gibbons-Hawking term, which is the surface integral of the extrinsic curvature, as well as from the bulk part. In the procedure, the extrinsic curvature term is canceled by a term extracted from the bulk part, and the divergent structure of the BAV in (7) is determined by the terms originated from bulk part that are expressed in terms of the intrinsic boundary geometry.

On the other hand, it must be also noted that in the above procedure, we have not used the full Einstein’s equations in obtaining the counterterm action, but only the constraint equation, $\mathcal{H}_\rho = 0$, has been used. In the following, we shall show that in the case of simple boundary geometry (round sphere), others in Gauss-Codazzi equations become trivial. Moreover, in the case of nontrivial boundary geometry, this scheme leads us to somewhat interesting observation. It will be presented in section 4.

3 AdS Space and Holographic Anomaly

The counterterm action for asymptotic AdS spaces in (1) is useful for various boundary geometries. However, evaluation of counterterm actions for higher dimensional

boundaries is not manageable for its mathematical difficulty. In the expression of counterterm action given in (9), we consider a simple but important example, Euclidean AdS space with S^d boundary and obtain a general form of the counterterm action available for any d -dimensional boundary.

The Euclidean AdS space with S^d boundary is described by the line element

$$\hat{G}_{\mu\nu}dx^\mu dx^\nu = \left(1 + \frac{r^2}{\ell^2}\right)^{-1} dr^2 + r^2 d\Omega_d^2. \quad (10)$$

The functional A in Eq.(7) for the metric (10) becomes

$$\begin{aligned} A(x^a; r_0) &= \frac{1}{8\pi G} \int^{r_0} dr \sqrt{\gamma_d} r^d \left(1 + \frac{r^2}{\ell^2}\right)^{-1/2} \left(R + \frac{d(d-1)}{\ell^2}\right) \\ &= -\frac{(d(d-1))^{(d+2)/2}}{16\pi G \ell} \sqrt{\gamma_d} \int^{R_0} dR R^{-(d+2)/2} \left(1 + \frac{\ell^2 R}{d(d-1)}\right)^{1/2}, \end{aligned} \quad (11)$$

where R_0 denotes the scalar curvature on the boundary and γ_d is the metric of d -dimensional unit sphere. In the second line of Eq.(11), $d(d-1)/r^2 = R$ was used. After some algebraic calculation, we obtain

$$\begin{aligned} A(x^a; r_0) &= -\frac{(d(d-1))^{(d+2)/2}}{16\pi G \ell} \sqrt{\gamma_d} \left(\frac{2}{d(d-1)} \sqrt{1 + \frac{\ell^2 R}{d(d-1)}} \right. \\ &\quad \times \left\{ -\frac{d-1}{R^{d/2}} + \sum_{k=1}^{(d-2)/2} \left[\left(-\frac{\ell^2}{d(d-1)}\right)^k \prod_{m=1}^k \left(\frac{d-2m+1}{d-2m}\right) R^{-(d-2k)/2} \right] \right\} \\ &\quad \left. - \frac{1}{d} \left(-\frac{\ell^2}{d(d-1)}\right)^{d/2} \prod_{k=1}^{(d-2)/2} \left(\frac{d-2k-1}{d-2k}\right) \ln \frac{\sqrt{1 + \ell^2 R/(d(d-1))} - 1}{\sqrt{1 + \ell^2 R/(d(d-1))} + 1} \right) \end{aligned} \quad (12)$$

in even of d and

$$\begin{aligned} A(x^a; r_0) &= \frac{d(d(d-1))^{d/2}}{8\pi G \ell} \sqrt{\gamma_d} \left(1 + \frac{\ell^2 R}{d(d-1)}\right)^{3/2} \\ &\quad \times \left(-\frac{\ell^2}{d(d-1)}\right)^{(d-5)/2} \sum_{k=0}^{(d-3)/2} \prod_{m=0}^k \left(\frac{d-2m-1}{d-2m}\right) \end{aligned} \quad (13)$$

in odd of d . (After the Eq.(12), we dropped the subscript ‘0’ of the scalar curvature for simplicity.) Then, an *arbitrary* dimensional counterterm action for AdS spaces with S^d boundary is given by a polynomial in the boundary scalar curvature R as follows

$$\tilde{S} = \frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{g_0} \left(\frac{d-1}{\ell} + \frac{\ell}{2(d-2)} R - \frac{\ell^3}{8d(d-1)(d-4)} R^2 \right)$$

$$+\frac{\ell^5}{16(d(d-1))^2(d-6)}R^3+\cdots\Big), \quad (14)$$

where the terms in the parenthesis of Eq.(14) are terminated by

$$\frac{1}{2}(-1)^{(d+2)/2}\prod_{k=1}^{(d+2)/2}\left(\frac{2k-3}{2k}\right)\frac{\ell^{d+1}}{(d(d-1))^{d/2}}R^{(d+2)/2}, \quad (15)$$

in the case of $d = \text{even}$, and

$$(-1)^{(d+1)/2}\prod_{k=1}^{(d-1)/2}\left(\frac{2k-3}{2k}\right)\frac{\ell^{d-2}}{(d(d-1))^{(d-3)/2}}R^{(d-1)/2}, \quad (16)$$

in odd d case. Using a relation $R_{ab}R^{ab} = R^2/d$, it can be shown that the counterterm action in (14) is equivalent to the Eq.(1). That is, the counterterm action in (1) can be written by a polynomial in the boundary scalar curvature R terminated by the terms given in (15) or (16) for AdS spaces with S^d boundary.

On the other hand, the counterterm action for even dimensional boundary in (14) fails on eliminating all divergent terms appearing in the BAV. Instead, the RA contains a logarithmic divergent term

$$\frac{1}{16\pi G}\int d^d x \sqrt{g_0}(-1)^{d/2}\prod_{k=1}^{d/2}\left(\frac{2k-3}{2k}\right)\frac{\ell^{d-1}}{(d(d-1))^{(d-2)/2}}R^{d/2}\ln R. \quad (17)$$

It has been already understood in the context of the AdS/CFT correspondence [6][11]; The regularization of BAV by introducing local counterterms may break conformal invariance and RA is left with a logarithmic divergent term. According to this prescription, considering a scale transformation $\delta r = r\delta\epsilon$ for an infinitesimal constant parameter $\delta\epsilon$, the holographic conformal anomaly, $\mathcal{A}$, for which dual CFT is coupled to the background gravity with S^d boundary is given by

$$\mathcal{A} = -\frac{1}{8\pi G}(-1)^{d/2}\prod_{k=1}^{d/2}\left(\frac{2k-3}{2k}\right)\frac{\ell^{d-1}}{(d(d-1))^{(d-2)/2}}R^{d/2}. \quad (18)$$

The conformal anomaly in arbitrary dimensions has been given in geometric description [19]. Restricting the CFT in background S^d geometry, Eq.(18) is an alternative expression of the conformal anomaly in arbitrary dimensions.

For S^2 boundary, the Eq.(18) recovers well known result

$$\mathcal{A}_{d=2} = -\frac{\ell}{16\pi G}R. \quad (19)$$

Comparing the $(1+1)$ -dimensional anomaly on a surface of radius ℓ , $-1/(8\pi G\ell) = -c/(12\pi\ell^2)$, the central charge c becomes $3\ell/(2G)$. From the Eq.(18) for $d = 4$, we find that the conformal anomaly agrees with that of Ref.[11]

$$\mathcal{A}_{d=4} = \frac{\ell^3}{768\pi G} R^2 = \frac{\ell^3}{8\pi G} \left(-\frac{1}{8} R_{ab} R^{ab} + \frac{1}{24} R^2 \right). \quad (20)$$

The conformal anomaly for $\mathcal{N} = 4$ super Yang-Mill theory on S^4 is $3N^2/(8\pi^2\ell^4)$. Comparing with the anomaly on this boundary from the Eq.(20), $3/(16\pi G\ell)$, we obtain the expected result

$$N^2 = \frac{\pi\ell^3}{2G}, \quad (21)$$

where N is the rank of the gauge group of the dual $\mathcal{N} = 4$ supersymmetric $d = 4$ $SU(N)$ YM theory. At last, it can be seen that for six dimensional boundary, the anomaly in (18) is equivalent to that given in [11]

$$\begin{aligned} \mathcal{A}_{d=6} &= -\frac{\ell^5}{115200\pi G} R^3 \\ &= -\frac{1}{16\pi G} \left(\frac{\ell^5}{64} \right) \left(\frac{1}{2} R R_{ab} R^{ab} - \frac{3}{15} R^3 - R^{ab} R_{acbd} R^{cd} \right. \\ &\quad \left. + \frac{1}{5} R^{ab} D_a D_b R - \frac{1}{2} R^{ab} \square R_{ab} + \frac{1}{20} R \square R \right). \end{aligned} \quad (22)$$

In fact, since we are concerned about S^6 boundary, the terms in third line including derivatives vanish. On the other hand, Eq.(22) can be verified by considering the central charge of N coincident M5-branes in the large N limit. It has been shown that the central charge is proportional to N^3 [20]. So, the anomaly on S^6 boundary with radius ℓ , $15/(64\pi G\ell)$, is proportional to $N^3/(\pi^4\ell^6)$. Thus, we find [5]

$$N^3 \sim \frac{\pi^3\ell^5}{G}. \quad (23)$$

Before ending of this section, it is useful on the next section to consider another Euclidean AdS space with different boundary geometry, $S^{d-1} \times S^1$,

$$ds^2 = \left(1 + \frac{r^2}{\ell^2} \right) d\tau^2 + \left(1 + \frac{r^2}{\ell^2} \right)^{-1} dr^2 + r^2 d\Omega_{d-1}^2. \quad (24)$$

For the $S^{d-1} \times S^1$ boundary, the functional A in (7) becomes

$$A(x^a; r_0) = \frac{\sqrt{\gamma_{d-1}}}{8\pi G} \left(\frac{d-1}{\ell^2} \right) \left(\frac{(d-1)(d-2)}{R} \right)^{d/2} \left(1 + \frac{\ell^2 R}{(d-1)(d-2)} \right). \quad (25)$$

Since all terms in expanding of Eq.(25) are divergent for $d > 2$, the counterterm action is just the negative of S_{cl} in (7) [14]

$$\tilde{S} = \frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{g_0} \left(\frac{d-1}{\ell} \right) \left(1 + \frac{\ell^2 R}{(d-1)(d-2)} \right)^{1/2}. \quad (26)$$

It can be shown that using $R_{ab}R^{ab} = R^2/(d-1)$ and expanding for ℓ , the counterterm action in (26) is equivalent to Eq.(1). However, it must be noted that while the counterterm action for S^d boundary is given by a finite sum of the series in (1), for $S^{d-1} \times S^1$ boundary it is given by an infinite sum. As mentioned in Ref.[14], in the process the divergent factors $1/(d-4)$, $1/(d-6)$, $\dots$ in (1) are canceled. Thus, while conformal invariance of the RA for S^d boundary is broken by the anomaly in (18), for the $S^{d-1} \times S^1$ boundary it is still conformal invariant.

4 AF Space and Holographic Anomaly

Now, consider counterterm actions for asymptotic flat spaces. In Ref.[14], it has been shown that the counterterm action for AF spaces is not be able to obtain on taking a limit of $\ell \rightarrow \infty$ in the procedure adapted for AdS spaces, an iteration process using the Gauss-Codazzi equations, because of mathematical difficulty. However, in our procedure, those are simply obtained by taking the $\ell \rightarrow \infty$ limit on the functional A 's

$$\tilde{S} = -\frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g_0} \sqrt{\frac{dR}{d-1}} \quad (27)$$

in (12) and (13), and

$$\tilde{S} = -\frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g_0} \sqrt{\frac{d-1}{d-2}} R \quad (28)$$

in (25). In Eqs.(27) and (28), the counterterm actions were written in the Lorentzian signature. Given in Ref.[14], the counterterm actions in (27) and (28) can be written by

$$\tilde{S} = -\frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g_0} \sqrt{\frac{R^3}{R^2 - R_{ab}R^{ab}}}. \quad (29)$$

In fact, the expression of counterterm action for AF spaces in (29) is more general than those in (27) and (28). Because it is available for the AF spaces with $S^{d-n} \times \mathbb{R}^n$

boundary geometries described by the metric

$$\hat{G}_{\mu\nu}dx^\mu dx^\nu = (-dt^2 + dx_1^2 + \cdots + dx_{n-1}^2) + dr^2 + r^2 d\Omega_{d-n}^2. \quad (30)$$

For AF spaces described by the metric (30), the functional A becomes

$$A(x^a; \rho_0) = \frac{1}{8\pi G} \int^{\rho_0} d\rho N \sqrt{-g} R, \quad (31)$$

and then the counterterm action is

$$\begin{aligned} \tilde{S} &= -\frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g_0} \sqrt{\frac{d-n}{d-n-1}} R \\ &= -\frac{1}{8\pi G} \int_{\partial X} d^d x \sqrt{-g_0} \sqrt{\frac{R^3}{R^2 - R_{ab}R^{ab}}}. \end{aligned} \quad (32)$$

Up to now, we have considered counterterm actions for manifolds with simple boundary geometry (round sphere). Now, we speculate divergences of BAV due to deviation of boundary from the round sphere. In Ref.[14], an AF space in spheroidal coordinates given by

$$\hat{G}_{\mu\nu}dx^\mu dx^\nu = -dt^2 + \frac{\rho^2}{r^2 + a^2} dr^2 + \rho^2 d\theta^2 + \sin^2 \theta (r^2 + a^2) d\phi^2 + r^2 \cos^2 \theta d\Omega_{d-3}^2 \quad (33)$$

was investigated. This space can be obtained by setting the mass to zero in higher dimensional Kerr metric [18]

$$\begin{aligned} \hat{G}_{\mu\nu}dx^\mu dx^\nu &= -\frac{\Delta}{\rho^2} (dt - a \sin^2 \theta d\phi)^2 + \frac{\sin^2 \theta}{\rho^2} (adt - (r^2 + a^2) d\phi)^2 \\ &\quad + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + r^2 \cos^2 \theta d\Omega_{d-3}^2, \end{aligned} \quad (34)$$

where $\rho^2 = r^2 + a^2 \cos^2 \theta$, $\Delta = r^2 - 2mGr^{4-d} + a^2$, and m and a are the black hole mass and the angular momentum per unit mass, respectively. It is important that the metric in (33) does not describe the asymptotic spacetime of the Kerr black hole in (34). Because, in the process, one is to meet a naked singularity. It is just the flat spacetime metric of $n = 1$ in Eq.(30) written in spheroidal coordinates.

The functional $A(x^a; r_0)$ for the metric (33) becomes

$$\begin{aligned} A(x^a; r_0) &= \frac{\sqrt{\gamma_{d-3}}}{8\pi G} \int^{r_0} dr r^{d-3} \left[\frac{2a^2 ((d-3) \sin^2 \theta - \cos^2 \theta)}{r^2 + a^2 \cos^2 \theta} \right. \\ &\quad \left. + (d-1)(d-2) + (d-3)(d-4) \frac{a^2}{r^2} \right] \sin \theta \cos^{d-3} \theta \end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{\gamma_{d-3}}}{8\pi G} r_0^{d-2} \left[(d-1) + \left(d-3 + \frac{2((d-3)\sin^2\theta - \cos^2\theta)}{d-4} \right) \frac{a^2}{r_0^2} \right. \\
&\quad \left. - \frac{2\cos^2\theta((d-3)\sin^2\theta - \cos^2\theta)}{d-6} \frac{a^4}{r_0^4} + \dots \right] \sin\theta \cos^{d-3}\theta, \quad (35)
\end{aligned}$$

where the divergence terms in the bracket are terminated by

$$2a^2 ((d-3)\sin^2\theta - \cos^2\theta) (-a^2 \cos^2\theta)^{(d-4)/2} r_0^{-(d-2)} \ln r_0 \quad (36)$$

in even d , and

$$2a^2 ((d-3)\sin^2\theta - \cos^2\theta) (-a^2 \cos^2\theta)^{(d-5)/2} r_0^{-(d-3)} \quad (37)$$

in odd d , respectively. In the above calculation, the d -dimensional scalar curvature R is

$$\begin{aligned}
R &= \frac{2a^2 ((d-3)\sin^2\theta - \cos^2\theta)}{(r^2 + a^2 \cos^2\theta)^2} + \frac{(d-3)(d-4)}{r^2 \cos^2\theta} \\
&\quad + \frac{(2(2d-5) - (d-3)(d-4)\tan^2\theta)}{r^2 + a^2 \cos^2\theta}. \quad (38)
\end{aligned}$$

The appearance of logarithmic divergence in Eq.(36) looks like strange. But, this is an artificial effect due to the fact that we have not used yet the remaining Einstein's equations that are normal-tangential and tangential-tangential projections. In fact, rewriting the terms in the bracket of Eq.(35) as

$$\begin{aligned}
&(d-1) + \left(d-2 - \cos^2\theta + \frac{d\sin^2\theta - 2}{d-4} \right) \frac{a^2}{r_0^2} \\
&+ \left((\cos^4\theta - \cos^2\theta) + \frac{\cos^2\theta(\cos^2\theta - d\sin^2\theta)}{d-6} \right) \frac{a^4}{r_0^4} + \dots, \quad (39)
\end{aligned}$$

comparing with the calculation of Ref.[14], some additional terms appear, $r_0^{-2}/(d-4)$, $r_0^{-4}/(d-6)$, $\dots$, which may cause the logarithmic divergent term of (36). After substituting the remaining equations into the action, those additional terms and the logarithmic divergence would vanish.

However, this logarithmic divergence seems to tell us something more. Following the procedure taken in the AdS descriptions, the logarithmic divergence breaks conformal invariance of the RA within a conformal anomaly

$$\mathcal{A}^{flat} = \frac{r_0^{d-1}}{4\pi G} \left[\frac{a^2 (-a^2 \cos^2\theta)^{(d-4)/2} ((d-3)\sin^2\theta - \cos^2\theta)}{r_0^{2(d-2)} \sqrt{(r_0^2 + a^2 \cos^2\theta)(r_0^2 + a^2)}} \right]. \quad (40)$$

For a strict argument, consider 4-dimensional boundary whose the conformal anomaly, $\mathcal{A}_{d=4}^{flat}$, is

$$\mathcal{A}_{d=4}^{flat} = \frac{r_0^3}{4\pi G} (\sin^2 \theta - \cos^2 \theta) \left(\frac{a^2}{r_0^6} \right) \left(1 + \frac{a^2 \cos^2 \theta}{r_0^2} \right)^{-1/2} \left(1 + \frac{a^2}{r_0^2} \right)^{-1/2}. \quad (41)$$

Then we find that up to leading order the anomaly is proportional to $\square R$

$$\mathcal{A}_{d=4}^{flat} \sim \frac{r_0^3}{4\pi G} \left(-\frac{1}{40} \square R + \mathcal{O} \left(\frac{a^4}{r_0^8} \right) \right). \quad (42)$$

As well known in the dual field theory on curved boundary, the 4-dimensional anomaly has an ambiguity that $\square R$ term can be added to the anomaly with an undetermined coefficient. This corresponds to the choice of different schemes for regularizing the field theory. In our case, this comes under different choices for the counterterm action. So, we can add additional counterterms given by

$$\Delta \tilde{S} \sim r_0^3 \int_{\partial X} d^4 x \sqrt{-g_0} (aE + bC_{abcd}C^{abcd} + cR^2), \quad (43)$$

where E is the Euler invariant $E = R_{abcd}R^{abcd} - 4R_{ab}R^{ab} + R^2$ and C^{abcd} is the Weyl tensor. Taking the coefficients as $b = -a$, $c = 0$, then the additional counterterms become

$$\Delta \tilde{S} \sim -2ar_0^3 \int_{\partial X} d^4 x \sqrt{-g_0} \left(R_{ab}R^{ab} - \frac{1}{3}R^2 \right). \quad (44)$$

The additional counterterm action in (44) is just that observed by Solodukhin in Ref.[17]; Those counterterms cancel the divergent terms which are caused by deviation of boundary from round sphere. It should be also noted that in spite that the counterterm actions in Eqs.(28) and (29) based on round sphere boundary exactly cancel divergent terms of the gravitational action for $d < 6$, the additional counterterm action in (44) is not proportional to the cubed boundary curvature $r_0^3 \int_{\partial X} R^3$ but $r_0^3 \int_{\partial X} R^2$. This is because the first correction (i.e., small deviation from round sphere) identically vanishes [17], i.e., leading terms of $R_{ab}R^{ab}$ and R^2 are canceled each other in the brace of Eq.(44).

5 Summary and Discussions

The counterterm subtracting method to define a finite gravitational action on non-compact spacetime has been speculated. It has been shown that using the ADM

formalism, the counterterm action could be explicitly written in terms of the intrinsic boundary geometry. On the other hand, using the form of counterterm action, we have obtained an expression of counterterm action available for *arbitrary* dimensional AdS spaces with S^d boundary geometry. Moreover, from this expression the arbitrary dimensional conformal anomaly has been driven. Our additional observation is that the counterterm action for AF spaces can be obtained as taking the limit of $\ell \rightarrow \infty$ in the procedure adapted for AdS spaces.

Another interesting observation in the description for counterterm action developed in this paper has been given in the example of AF space with nontrivial boundary geometry. It has been shown that the additional counterterms to eliminate (leading) divergent terms due to deviation of boundary from round sphere, which was suggested by Solodukhin [17], can be imagined from appearance of logarithmic divergence in BAV and perspective of the corresponding anomaly proportional to $\square R$. It seems that it is due to a deceptive procedure as skipping over tangential-tangential and tangential-normal projections of Einstein's equations in calculation of BAV. In fact, simply using the full Einstein's equations, then we would obtain a BAV without the logarithmic divergent term. However, it appears that there is something in the holographic sense. For the 5-dimensional Kerr-AdS spacetime with the boundary of 4-dimensional rotating Einstein universe, the trace of stress tensor does not vanish, but the evaluated BAV does not contain a corresponding logarithmic divergence [21]. As mentioned in the Ref.[21], the corresponding logarithmic divergence of the BAV should not be present for a spacetime which can be written locally as a product. It has been also observed that the conformal anomaly, which corresponds to the CFT anomaly, is also proportional to the $\square R$ term. In addition, it can be shown [22] that taking the limit of $\ell \rightarrow \infty$ on the Kerr-AdS description, the conformal anomaly becomes the Eq.(42). These holographic similarities of the Kerr spacetime with the Kerr-AdS description will be studied in detail in Ref.[22]. It is also expected that this study will be of help to prescribe the problem of construction of the flat-space S-matrix which has been studied on the large radius limit in the AdS/CFT correspondence and suffered from non-local holographic mapping [23][24].

Acknowledgments: I thank V. Frolov, A. Zelnikov, and Y. Gusev for helpful

discussions. This work was supported in part by Korea Science and Engineering Foundation and in part by National Science and Engineering Research Council of Canada.

References

- [1] G.W. Gibbons and S.W. Hawking, Phys. Rev. D**15** (1977) 2752.
- [2] J.D. Brown and J.W. York Jr, Phys. Rev. D**47** (1993) 1407.
- [3] G.T. Horowitz and S.W. Hawking, Class. Quantum Grav. **13** (1996) 1487.
- [4] V. Balasubramanian and P. Kraus, Commun. Math. Phys. **208** (1999) 413.
- [5] J. Maldacena, Adv. Theor. Math. Phys. **02** (1998) 231.
- [6] E. Witten, Adv. Theor. Math. Phys. **02** (1998) 253.
- [7] S.S. Gubser, I.R. Klebanov, and A.M. Polyakov, Phys. Lett. B**428** (1998) 105.
- [8] G. 't Hooft, *'Dimensional Reduction in Quantum Gravity'*, in *Salamfest* (1993) p.284, gr-qc/9310026.
- [9] L. Susskind, J. Math. Phys. **36** (1995) 6377.
- [10] L. Susskind and E. Witten, *'The Holographic Bound in Anti-de Sitter Space'*, hep-th/9805114.
- [11] M. Henningson and K. Skenderis, JHEP **07** (1998) 023.
- [12] S. Hyun, W.T. Kim, and J. Lee, Phys. Rev. D**59** (1999) 084020.
- [13] R. Emparan, C.V. Johnson, and R.C. Myers, Phys. Rev. D**60** (1999) 104001.
- [14] P. Kraus, F. Larsen, and R. Siebelink, Nucl. Phys. B**563** (1999) 259.
- [15] N.D. Birrell and P.C.W. Davies, *'Quantum Fields in Curved Space'*, Cambridge University Press, Cambridge (1982).
- [16] S. Nojiri and S.D. Odintsov, Phys. Lett. B**444** (1998) 92.

- [17] S.N. Solodukhin, ‘*How to make the gravitational action on non-compact space finite*’, hep-th/9909197.
- [18] R.C. Myers and M.J. Perry, Ann. Phys. **172** (1986) 304.
- [19] S. Deser and A. Schwimmer, Phys. Lett. B**309** (1993) 279.
- [20] S.S. Gubser and I.R. Klebanov, Phys. Lett. B**413** (1997) 41.
- [21] Adel M. Awad and Clifford V. Johnson, *Holographic Stress Tensors for Kerr-AdS Black Holes*, hep-th/9910040.
- [22] Jeongwon Ho, *in preparation*.
- [23] L. Susskind, *Holography in the Flat Space Limit*’, hep-th/9901079.
- [24] S.B. Giddings, *Flat-Space Scattering and Bulk Locality in the AdS/CFT Correspondence*’, hep-th/9907129.